

PATENT 0425-0847P

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicant:

Koichi ITO et al.

Conf.:

9635

Appl. No.:

09/913.444

Group:

1624

Filed:

August 15, 2001


Examiner: Bruck KIFLE

For:

HETERODIAZINON COMPOUND

DECLARATION UNDER 37 C.F.R. \$ 1.132

. I, Takahisa Hanada, declare as follows:



- 1. I am one of the co-inventors of the invention claimed in the above-referenced application. I have carried out comparative tests which are fully described herein, to support enablement of the invention, and to show unexpected superior properties of the compounds of the present invention over U.S. Patent 4,670,555 to Dekeyser et al.
- 2. A Formalin assay was conducted either by myself or under my supervision and control to show that the present invention is effective in treating pain. In the assay, the analgesic effect of the compound of claim 1, particularly compound #54 was tested. In the test, mice were anesthetized lightly with halothane. 20 µl of 3% formalin solution was injected subcutaneously into the dorsal

surface of the left hind paw of the mice. The mice were them placed in an observation chamber. Compound #54 was intravenously administered to the mice just before the formalin injection. The mice demonstrated pain related licking and biting behavior. These responses were measured during the first five minutes after the formalin injection. The example Compound #54 reduced the pain related licking and biting behavior in the mice. This result indicated that compound #54 has an analgesic effect. The results are shown in Table 1 attached hereto.

3. An BAB assay was conducted with Lewis rats. The Lewis rats were immunized s.c. in each hind foot with 25 µl of inoculum containing 50 µl MBP emplsified in Freund's complete adjuvant containing M. tuberculosis. Thereafter, a vehicle (methyl cellulose (MC)) or example compound #1 was administered to the rats. On day 7, post immunization, 150 and 3000 mg/kg of each were administered twice daily p.o. This administration continued up to and including day in body weight and post-immuization. Changes neurological scores were monitored everyday. The results are demonstrated in Table 2 attached hereto. Compound #1 was shown to be effective in eliminating paralysis and an unstable gait in the rats. The results indicate that Compound #1 is effective in treating epilepsy.

Comparative in vitro and in vivo tests were conducted utilizing example compounds 1, 33, 34, 42, 54, 61, 66 and 68 listed in the specification. The in vitro assay was conducted in the same manner as in the specification at pages 107-108. The results of the in vitro assay are reported in Table 3 attached hereto. Examples 2, 5, 6, 8, 14, 27, 28 and 29 of U.S. Patent 4,670,555 to Dekeyser et al. were also tested in the same manner. The results of the in vitro assay are also reported in Table 3. The comparative results demonstrate that the IC50 values for the compounds of the present invention are unexpectedly superior to the compounds in Dekeyser.


An in vivo assay was conducted to test the abovementioned compounds of the invention against the abovementioned compounds of Dekeyser. The in vivo assay was conducted in the same manner as the in vivo assay described in the specification at pages 112-113. The action of the test compound was judged by the presence or absence of the mice. A total of six mice were seizures 1/29/2004 T.H. tested. The results are reported in Table 3 in terms of the number of mice out of 6, that actually had aparens. seizures The results of the in vivo assay demonstrate that the compounds of the present invention are more effective in reducing epasses than the compounds of Dekeyser et al. Scizures 1/19/2004 T.M These results were unexpected.


1/29/2004 T-H. own knowledge are true. Statements made on information and belief are believed to be true to the best of my knowledge. These statements were made with the full knowledge that willfully providing false statements and the like are punishable by fine and/or imprisonment under \$1001 of Title 18 of the United States Code, and that willful false statements may jeopardize the validity of the application or any patent that may grant therefrom.

DATED: 1/29/2004 BY	Y: Takahisa Hanada Takahisa HANADA
---------------------	---------------------------------------

Attachments: Tables 1-3


TABLE 1











The following wording system was used to grade neurological impairment:

Normal 0;

Half paredyzed tad Ó.B;

Fraccied tall 1:

Shently materity con 15

प्रिकास स्वीतास्त्रात्म Z;

Complete penalysis of one hindlind 2.57

Complete persivels of both hindlimbs 87

Complete parelycle of both hindlimbs 8.5

Complete paralysis of both bindinals


4:

and workness of one firelimb and weatness of both forelimbs

Mariband/death. 5;

Results

Effect of example #1 compound on disease symptom.



FEB 1 0 2004 E

TABLE 3



						4 m - 10 m m		
Citad	reference(US 4	L870.555)			Our invention			
alamex	Formula	ו פדמאי אוו	מיזיי חו	Ecample No.	Fermula	CHN	0 10	y îvo
Vo. 2	90	3.94		1	0.00	a .	11 <u>.</u> 28 (4	/6 2)
5	چ گڻ	18.9	5/6	33	. 9.		4.28 Q	/6
8	200	10.4	46/5	34	Q Q	g F	9.21 0	√s
	S.O.	○ 18 <u></u>	4/4	4	25		3.78)/G
1	S. C.		66/6	5	4	\$. }	3.54	1/6
	33		02 6/6		725	5	4.61	0/6_
		Q	17 6/6		~ \$ \\ \frac{1}{2} \\	3	272	1/8
	29		16 5/6		58 The value	}		1/6_

In vitro:Pharmacological experimental example 1. The value indicates IC50
In vitro:Pharmacological experimental example 8.A text compound 30mg/kg was
administered to each mouse. The value indicates the number of epsemed mine in 6
mice.

(#1): I Simplifies of the compound was administered to each mouse.

(42). Thrus/Ice of the compound was administered to each mouse.